



Nitrogen Efficiency and Management

Issued September 2007

Nutrient Management Technical Note No. 3 was prepared by John Davis, Acting National Nutrient Management Specialist, Ecological Sciences Division, located at the NRCS National Headquarters, Washington, DC. For further information, contact the National Agronomist, 202-720-3783.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202)720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, SW., Washington, DC 20250-9410, or call (800)795-3272 (voice) or (202)720-6382 (TDD). USDA is an equal opportunity provider and employer.

Nitrogen Efficiency and Management

Overview

Nitrogen (N) is an essential element for plant growth and animal nutrition and is the nutrient taken up in the largest amount by plants. N is a component of essential plant compounds required in numerous biological processes, including genetic transmission (DNA, RNA), plant growth (proteins, enzymes) and photosynthesis (chlorophyll). Most non-leguminous crops require added N to achieve the yields expected in modern agriculture. Nitrogen is also the nutrient that is most likely to be lost to the environment.

Excess N in the environment can adversely affect humans, animals, and the biology of the environment. The movement of N into surface waters often leads to its contamination and eutrophication, a general enrichment in fertility, which later produces a decline in oxygen content of the water. of Nlik.lyes) anronmeation results in subsequent death of fish and decline in aquatic life. Nitrogen as a gas (nitrous oxide) can degrade the ozone level and contribute to global warming. Ammonia losses from soil and plants contribute to off-site contamination and

nitrogen compounds as it passes across the soil surface. The highest potential for nitrogen runoff comes from surface applied fertilizers and manures, even if erosion is minimal. Under these conditions, N losses in runoff can be high, but they can vary depending on the amount of nitrogen applied and the soil surface conditions.

Ammonia volatilization

Ammonium ions (NH_4^+) in the soil solution exist in equilibrium with ammonia (NH_3) in the soil solution. There is a tendency for the equilibrium to favor conversion to ammonia because solution NH_3 is subject to gaseous losses to the atmosphere.

Soil pH and the concentration of ammonium (NH_4^+) in the soil solution are important factors affecting amount of ammonia (NH_3) volatilized. As soil pH increases, the fraction of soil-solution ammonium and NH_3 increases by an order of magnitude for every unit of pH above 6.0.

The following summarizes favorable conditions for NH_3 volatilization:

- Prominent on calcareous soils, especially as soil pH exceeds 7;
- Can be appreciable for neutral or alkaline soils as they dry out;
- Increases with temperature and wind speed;
- Greater in soils of low CEC, such as sands where NH_4^+ adsorption is low;
- High when high-N organic wastes, such as manure, are applied to the soil surface and permitted to decompose;
- High from urea when it is applied to grass or pasture, as a result of hydrolysis of the urea to NH_3 by urease enzyme;
- Losses are decreased by growing plants.

Volatilization losses can be as high as 100% of the ammonium on or near the soil surface if one or more of the above conditions exist.

Mineralization, Nitrification and Denitrification

As plants and other organic residues decompose, nitrogen is converted from organic-N into ammonium (NH_4^+), a process referred to as *mineralization*.

Nitrification

Is generally viewed as a two-step process involving several types of specialized bacteria.

Step 1. Ammonium (NH_4^+) \rightarrow Nitrite (NO_2^-)

In the presence of genus *Nitrosomonas* bacteria.

Step 2. Nitrite (NO_2^-) \rightarrow Nitrate (NO_3^-) occurs
In presence of genus *Nitrobacter* bacteria.

The over-all process results in the biological oxidation of ammonia with oxygen into nitrite followed with the oxidation of these nitrites into nitrates. Nitrification is an important step in the nitrogen cycle.

Denitrification

Nitrate can also be lost to the atmosphere through the *denitrification* processes. Nitrite (NO_2^-) usually does not accumulate in soils because it is rapidly transformed to NO_3^- or is denitrified to N_2 gas, nitrous oxide (N_2O), nitric oxide (NO), or one of the other gaseous nitrogen oxide (NO_x) compounds. Nitrous oxide, a product of incomplete denitrification, is a greenhouse gas and may contribute to global climate change and to thinning of the ozone layer.

Denitrification is the route for most losses of gaseous N compounds to the atmosphere. The potential for denitrification is increased as oxygen levels in the soil decrease. Nitrate, a desirable nutrient for plants, is reduced first to NO_2^- , then to NO , next to N_2O , and finally to N_2 . In addition to limited oxygen, denitrification needs a carbon source of energy, effective microbes, and oxides of N. Denitrification losses can be as high as 10 pounds of nitrogen per acre per day when conditions are optimum.

N Management Practices that Improve Nitrogen Efficiency

The objective of N management is to maximize the efficiency of plants to use the applied N. Improvements in nitrogen use efficiency are associated with decreases in N loss through processes of fate and transport mentioned above. While losses cannot be avoided completely, significant improvements can be realized by applying management and conservation practices to the cropping system. The following

list of techniques and practices provide some general guidance.

Conservation Tillage

- Conservation tillage reduces total N losses due to reduced sediment and runoff.
- Conservation tillage optimizes soil moisture conditions that improve water use efficiency.
- Runoff and leaching losses of nitrate are not consistently affected by conservation tillage.

Other Conservation Practices

- Crop rotations favorably impact the fertility of the soil. Deep-rooted crops capture nitrates lower in the soil profile.
- Cover crops capture residual nitrogen after crop harvest and recycle it as plant biomass.
- Soil erosion and runoff control lessens the nutrient transport processes.
- Incorporation and/or injection lessen the volatilization of ammonium nutrients.

Nitrogen Rates

- Maintain accurate yield records or use an alternate realistic method to determine reasonable yield expectations.
- Follow your Land Grant University's recommendation for nitrogen application.
- Take credit for N applied as manure, irrigation water, atmospheric deposition, and fixed by legumes in rotation.
- Use appropriate soil tests to determine residual soil, water and plant N.

Soil Testing for N

- Preplant soil tests provide information on the soil's supply power.
- Late spring or pre-side-dress N tests can determine if and how much additional N is needed.
- New soil test and sampling procedures, such as amino sugar tests, grid mapping, and real-time sensors will define N requirements.
- Post-harvest soil tests determine if N management the previous season was appropriate.

Application Precision of N

- New designs of manifolds on application equipment increases the uniformity of anhydrous ammonia distribution.
- Row applicators, such as injectors, which form a compacted soil layer and surface ridge, can reduce N losses.
- Variable rate applicators, combined with intensive soil or crop sampling, allow more precise and responsive application rates.

Irrigation Management of N

- Schedule irrigation based on soil moisture estimates and daily crop needs to improve irrigation efficiency and soil moisture use.
- Sprinkler irrigation systems apply water more uniformly and in lower amounts than furrow or basin irrigation systems.
- Furrow irrigation efficiency can be improved by adjusting set time, stream size, furrow length, watering every other row, or the use of surge valves.
- Alternate row irrigation and fertilization minimizes water contact with nutrients.
- Application of N fertilizer through irrigation systems facilitates supplying N when crop demands are greatest.
- Polyacrylamide (PAM) treatment during furrow irrigation reduces sediment and N losses.

Crop Testing for N

- Plant tissue tests can identify N deficiencies.
- Sensing variations in plant chlorophyll content facilitates variable rate N applications in-season.
- Post-black-layer corn stalk nitrate tests help to determine if N rates were low, optimal, or excessive in the previous crop, so that management changes can be made in following crops.

Timing of N Applications

- Apply N close to the time when crops can utilize it.
- Make side-dress N applications close to the time of most rapid N uptake.

- Split applications, involving more than one application, allow efficient use of applied N and reduce the risk of N loss to the environment.
- Fall N applications may be discouraged in some areas, but can be successful if ammonium fertilizers and manures are applied when soil temperatures are below 50° F.

N Fertilizer Application Method and Placement

- Injection or incorporation of urea or N solutions reduces volatility losses.
- In ridged crops, placing N fertilizers in a band in ridges makes N less susceptible to leaching.

N Forms, including Slow or Controlled Release Fertilizers and Inhibitors

- Slow or controlled release fertilizer delays the availability of nitrogen to the plant until a time that is more appropriate for plant uptake and can avoid periods of potentially risky fate and transport.
- Nitrification inhibitors maintain applied ammoniacal fertilizers in the ammonium form longer, reducing leaching and denitrification losses.
- Where fall N applications are appropriate, nitrification inhibitors reduce risk of leaching loss.
- Urease inhibitors temporarily block the function of the urease enzyme, maintaining urea-based fertilizers in the non-volatile urea form, reducing volatilization losses when these fertilizers are surface applied in high residue, conservation tillage systems.

Drainage Tile Systems

- Subirrigation systems recycle nitrate leached from the soil profile and reduce nitrate lost to subsurface tile drainage.
- Controlled drainage favors denitrification.

Conservation Buffers

- Buffers trap sediment containing ammonia and organic N.
- Nitrate in subsurface flow is reduced through denitrification enhanced by carbon energy sources contained in the soil associated with buffer vegetation.
- Buffer vegetation takes up nitrogen, other nutrients, and reduces loss to water.

Constructed Wetlands

- Constructed wetlands located strategically on the landscape to process drainage effluent reduces sediment and nitrate loads to surface water.

Pest Management

- Appropriate pest management allows crops to attain their potential yields, optimizing applied N thus reducing the excess N available for loss.
- Bt corn prevents European corn borer feeding and associated stalk rots, which can cause corn to die early and leave excess N in the soil.

Breeding Crops for Efficient N Uptake

- Particular crop varieties are able to more efficiently extract N from the soil and improve N use efficiency.

References

- 1) Follett, R.F. 2001. Nitrogen Transformation and Transport Processes. pp. 17-44, In R.F. Follett and J. Hatfield. (eds.). 2001. Nitrogen in the Environment; Sources, Problems, and Solutions. Elsevier Science Publishers. The Netherlands. 520 pp.
- 2) van Es, H.M., S.D. Klausner, and W.S. Reid. 1991. Nitrogen and the Environment. Cornell Coop. Ser. Bull. 218. Cornell University, Ithaca, NY.